

08/962,721

**INTERMEDIATE RELEASE NICOTINIC ACID COMPOSITIONS
FOR TREATING HYPERLIPIDEMIA WHICH EXHIBIT AN
IN VIVO STAIR-STEPPED ABSORPTION CURVE**

Abstract

5 Intermediate release nicotinic acid formulations exhibiting an *in vivo* stair-stepped or
sigmoidally-shaped absorption curves when the plasma nicotinic acid or nicotinuric acid curves are
deconvoluted using the Wagner-Nelson method, which are suitable for oral administration once-a-day
preferably during the evening or at night for treating hyperlipidemia without causing drug-induced
hepatotoxicity to such a level that requires the therapy to be discontinued, are disclosed. The stair-
10 stepped or sigmoidal-shaped absorption profiles for nicotinic acid or nicotinuric acid absorbed from
the nicotinic acid formulations of the instant invention are characterized by three phases and by the
fact that significant quantities of nicotinic acid are absorbed during the first two phases. The first
phase generally ends at about 2.3 hours following ingestion and the second phase generally ends at
about 7.3 hours after consumption. Approximately 6% of the nicotinic acid dose is generally
15 absorbed by the end of the first phase and approximately 91% of the nicotinic acid dose is absorbed
by the end of the second phase.

009544-1-103497

1